

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 970 615 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

12.01.2000 Bulletin 2000/02

(51) Int. Cl.⁷: **A23D 7/00**, A23D 9/00,
A23G 3/00, A23C 13/14,
A23L 2/38, A61K 31/23

(21) Application number: 98905789.8

(22) Date of filing: 06.03.1998

(86) International application number:
PCT/JP98/00926

(87) International publication number:
WO 98/39977 (17.09.1998 Gazette 1998/37)

(84) Designated Contracting States:
DE FR GB

(30) Priority: 12.03.1997 JP 5779397

(71) Applicant: Kao Corporation
Tokyo 103-0025 (JP)

(72) Inventors:
• MORI, Hideki,
Kao Corporation, Research Lab.
Ibaraki 314-0103 (JP)

• WATANABE, Takuya,
Kao Corporation, Research Lab.
Ibaraki 314-0103 (JP)

(74) Representative: HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(54) FOOD CONTAINING FAT OR OIL

(57) The invention provides foods containing oil and/or fat which upon ingestion into the body, are hardly accumulated as body fat, are utilizable more positively as energy, contribute to the storage of glycogen in the liver and muscle to achieve effects such as recovery from fatigue, supplement of nutrition and increase of stamina at the time of exhaustion of physical strength, at the time of fatigue and at the time of exercise, are excellent in the oxidation stability of oils and/or fats contained therein, and are excellent in eating-feeling such as in dissolution in the mouth. The foods of the invention comprise at least 0.5 % by weight of a partial glyceride derived from a C₂₋₁₀ aliphatic acid.

EP 0 970 615 A1

Description

TECHNICAL FIELD

5 [0001] The present invention relates to a food containing oil and/or fat. In particular, the present invention provides foods, though which are ingested, which are hardly accumulated as body fat, are easily utilizable as energy, and contribute to the storage of stamina at the time of fatigue. Further, the present invention provides foods with excellence in the oxidation stability of oils and/or fats contained therein and excellent in eating-feeling such as dissolution in the mouth.

10 BACKGROUND ART

[0002] Humans take in a source of necessary energy mainly from sugars and lipids, but have surplus calories in the modern dietary habits under the present circumstances. In particular, lipids have high calorie value and are regarded as a factor for facilitating obesity upon surplus intake to cause adult diseases. However, there are few lipids which are effective in inhibiting the accumulation of body fat and in more positive utilization as energy and are excellent in stability and eating-feeling.

[0003] JP-A 8-60180 describes that a fat composition comprising diglycerides containing middle- and long-chain aliphatic groups has a weak but inhibitory effect on accumulation of body fat.

20 [0004] JP-A 5-168142 discloses a diglyceride containing two long-chain aliphatic groups and utilizes it as a bloom inhibitor for chocolate.

DISCLOSURE OF THE INVENTION

25 [0005] Focusing on the characteristics in metabolism of nutrition on oils and/or fats, the present inventors made extensive study for the purpose of providing foods containing oils and/or fats, in the case of which are ingested into the body, are hardly accumulated as body fat, are utilizable more positively as energy, contribute to the storage of glycogen in the liver and muscle to achieve effects such as recovery from fatigue, supplement of nutrition and increase of stamina at the time of exhaustion of physical strength, at the time of fatigue and at the time of exercise, are excellent in the oxidation stability of oils and/or fats contained therein, and are excellent in eating-feeling such as in dissolution in the mouth. As a result, the present inventors found that partial glycerides derived from specific middle- and short-chain aliphatic acids are advantageous oils and/or fats in that they are easily utilizable as energy as described above, their accumulation as body fat can be easily controlled, and further they are excellent in eating-feeling and oxidation stability, and the present invention was thereby completed.

35 [0006] The present invention relates to foods comprising at least 0.5 % by weight of a partial glyceride derived from a C₂₋₁₀ aliphatic acid. The number of carbons of the aliphatic acid is preferably 8 to 10.

[0007] The partial glyceride is preferably a diglyceride. More preferably, at least one of the two aliphatic acids constituting the diglyceride is a C₈₋₁₀ aliphatic acid, or at least one of two aliphatic acids constituting the diglyceride is a C₁₀ aliphatic acid, or both of the two aliphatic acids constituting the diglyceride are C₈₋₁₀ aliphatic acids. The partial glyceride is preferably capric acid diglyceride.

40 [0008] The foods preferably comprise 0.5 to 85 % by weight of the partial glyceride.

[0009] Other preferable foods are those wherein the partial glyceride is a diglyceride, at least one of two aliphatic acids constituting the diglyceride is a C₁₀ aliphatic acid, and at least 4.0 % by weight of this partial glyceride is contained.

45 [0010] The foods are preferably in the form of drinks or oral administration agent.

[0011] Further, the present invention relates to foods having the whole or a part of oils and/or fats in the fat-containing foods replaced by a partial glyceride, wherein this partial glyceride is derived from a C₂₋₁₀ aliphatic acid and the content of this partial glyceride in foods is particularly at least 0.5 % by weight for oral administration.

50 [0012] Those who take in the foods of the present invention are supplemented with energy. Further, accumulation of body fat is inhibited in those who have taken in the foods. The partial glyceride is easily consumed as energy, and hardly accumulated in internal organs and adipose tissues. That is, glycogen is not consumed and readily stored. Accordingly, storage of glycogen is improved. This effect is more significant particularly, if at least 0.5 % by weight of oil and/or fat contained in the foods is substituted by the partial glyceride.

[0013] When the foods of the present invention are ingested, as compared with no ingestion of the foods, fats other than glycogen are predominantly consumed thereby significantly reducing the decrease of glycogen.

55 [0014] In the view of the significant effect, besides the above-mentioned action and effect, it is suggested that the partial glyceride may inhibit the accumulation of other oils and/or fats (adiposes) in internal organs or adipose tissues, or may promote the consumption of accumulated fats.

DETAILED DESCRIPTION OF THE INVENTION

[0015] Hereinafter, the present invention is described in detail.

[0016] In the present invention, the partial glyceride includes monoglycerides and diglycerides. However, C₂₋₁₀ monoglycerides usually taste bitter etc. It may be necessary, when the C₂₋₁₀ monoglyceride is used as foods, that they should be formed into foods making use of the flavor or that they should be used in combination with a masking agent for flavor. On the other hand, diglycerides taste less bitter, so they can be applied to a wide variety of foods, and for this reason, the foods preferably contain a large amount of diglycerides rather than a large amount of monoglycerides.

[0017] The aliphatic acid mentioned above is a C₂₋₁₀ aliphatic acids such as acetic acid, butyric acid, caproic acid, caprylic acid and capric acid. If the number of carbon atoms therein is low, peculiar flavor tends to be generated. Therefore the peculiar flavor needs a masking agent or needs a suitable foods for its unique flavor. And then, C₈₋₁₀ aliphatic acids with weak flavor are more preferable because of application to a wide variety of foods. Particularly, one of the two aliphatic acids of the diglyceride is preferably a C₈₋₁₀ aliphatic acid, especially preferably a C₁₀ aliphatic acid, and both of them are more preferably C₈₋₁₀ aliphatic acids and most preferably C₁₀ capric acid diglyceride.

[0018] The foods referred to in the present invention are the modes of foods containing oils and/or fats, which are ready to be eaten. The content of the partial glyceride in the foods is 0.5 % by weight or more, preferably 1 % by weight or more, more preferably 2 % by weight or more, most preferably 4 % by weight or more, as compared with the foods. That is, comparatively high calorie foods containing at least 0.5 % by weight of oils and/or fats are significant in the effect of the present invention.

[0019] Ingredients other than the partial glyceride used in the present foods may be any ingredients being usable for food.

[0020] Examples of the present foods include baked confectionery such as cookies and biscuits, fried cakes such as potato chips, cakes, ice creams, meats such as sausages and hamburgers, stews, fried foods such as fries and tempuras, and drinks such as juices and nutrition drinks. Further, the foods may be creaming powder etc. insofar as the foods are placed in coffee or tea and contained in a necessary amount. Similarly, they may be dressings and mayonnaise on vegetables etc. satisfying a necessary amount in use, or margarine and fat spreads on breads etc. satisfying a necessary amount in use.

[0021] The foods of the present invention, into which the partial glyceride has been incorporated, are hardly accumulated as body fat and are easily converted into energy very rapidly. Further, it is believed that because this action is preferentially brought about, the foods also contribute to e.g. the maintenance of stamina by keeping glycogen in the liver and muscle. Accordingly, the foods are very effective in supplement of nutrition, recovery from fatigue and increase of stamina at the time of fatigue, at the time of reduction of physical strength, and before, during and after exercise.

[0022] The foods of the present invention are preferably in the form of an oral administration. The preparation form includes e.g. tablets, capsules, powder, granules and syrups. For their manufacturing, conventional excipients, disintegrators, binders, lubricants, pigments, diluents and so on are used. The excipients include glucose, lactose etc.; the disintegrators include starch, sodium alginate etc.; the lubricants include magnesium stearate, paraffin sulfate, talk etc.; and the binders include dimethylcellulose, gelatin, polyvinyl pyrrolidone etc.

EXAMPLES

[0023] Hereinafter, the effect of the present invention is described in more detail with reference to the Examples, which however are not intended to limit the present invention. The term "%" in the Examples refer to "% by weight" unless otherwise specified.

[Preparation of oils and/or fats]

[0024] Capric acid and glycerol were mixed at a molar ratio of 2 : 1 in the presence of a catalyst, that is, a commercial lipase preparation of immobilized 1,3-position-selective lipase (Lipoyme 3A™ provided by Novo Industry A. S.), and the mixture was reacted at 40 °C under reduced pressure. The lipase preparation was filtered out, and the final product was subjected to molecular distillation and purified for discoloration, deodorization etc. to give an oil and/or fat consisting of 85 % diglyceride and 15 % triglyceride (oil and/or fat preparation A).

[0025] Rapeseed oil-decomposed aliphatic acids were used and treated in the same manner as above to give an oil and/or fat consisting of 86 % diglyceride and 14 % triglyceride (oil and/or fat preparation B).

[0026] In addition, 1 mole of capric acid was added to 1 mole of oleic acid monoglyceride (O-95R provided by Kao Corporation) and was reacted at 60 °C under reduced pressure in the presence of the above lipase preparation as the catalyst. After the lipase preparation was filtered out, the final reaction product was subjected to molecular distillation, and the monoglyceride, aliphatic acid, capric acid-capric acid diglyceride components were removed as much as possible and then purified for discoloration and deodorization to give a fat consisting of 80.9 % diglyceride (0.5 % capric

acid-capric acid diglyceride, 50 % capric acid-oleic acid diglyceride, and 30 % oleic acid-oleic acid diglyceride) and 19.1 % triglyceride (oil and/or fat preparation C).

[0027] Further, capric acid, palmitic acid and glycerol were mixed at a molar ratio of 1 : 1 : 1 and reacted at 60 °C under reduced pressure in the presence of the above lipase preparation as the catalyst. After the lipase preparation was filtered out, the final reaction product was subjected to molecular distillation, and the monoglyceride, aliphatic acid, capric acid-capric acid diglyceride components were removed as much as possible and purified for discoloration and deodorization to give a fat consisting of 80.8 % diglyceride (0.5 % capric acid-capric acid diglyceride, 51 % capric acid-palmitic acid diglyceride, and 29 % palmitic acid-palmitic acid diglyceride) and 19.2 % triglyceride (oil and/or fat preparation D).

[0028] Further, capric acid, caproic acid and glycerol were mixed at a molar ratio of 1 : 1 : 1 and reacted at 60 °C under reduced pressure in the presence of the above lipase preparation as the catalyst. After the lipase preparation was filtered out, the final reaction product was subjected to molecular distillation and purified for discoloration and deodorization to give an oil and/or fat consisting of 1.1 % monoglyceride, 84.6 % diglyceride (26.5 % capric acid-capric acid diglyceride, 42.3 % capric acid-caproic acid diglyceride, and 15.8 % caproic acid-caproic acid diglyceride) and 14.3 % triglyceride (oil and/or fat preparation E).

[0029] The glyceride composition and aliphatic acid composition in each oil and/or fat were analyzed in the following manner.

• Measurement of glyceride composition distribution

[0030] After the oil and/or fat was silylated with a silylating agent (silylating agent TH provided by Kanto Kagaku Co., Ltd.). It was analyzed by gas chromatography with a hydrogen ion flame detector, equipped with a capillary column (e.g. DBTM-1, J&W Co., Ltd.), and the glyceride composition distribution in the oil and/or fat was determined from the retention time and peak area ratio in the obtained chart.

• Measurement of aliphatic acid composition

[0031] Analyzed by gas chromatography according to a method in "2.4.20.2-77 Process for Preparing Fatty Acid Methyl Esters" and "2.4.21.2-73 Composition of Aliphatic Acids" in "Standard Methods for the Analysis of Fats, Oil and Related Materials, compiled by Japan Oil Chemists' Society". The distribution of aliphatic acids was determined from the retention time and peak area ratio in the obtained chart.

[0032] The oil and/or fat compositions used are shown in Table 1.

Example 1

[0033] Oil and/or fat preparations A, B, C and D to which 600 ppm tocopherol was added were examined for their oxidation stability and solubilization in the mouth. Oxidation stability was examined at 120 °C according to "2.4.28 Stability Test toward Autoxidation, 2.4.28.2-93 CDM Test" in "Standard Test Methods for Oil and Fat Analysis, compiled by Japan Oil Chemical Association". The results are shown in Table 2.

[0034] From the results described above, the oil and/or fat preparation A was more excellent in oxidation stability than the oil and/or fat preparations B and C. Further, its dissolution in the mouth is excellent, so it was found that the oil and/or fat preparation A is a general-purpose oil and/or fat being easily applicable to a wide variety of foods.

Example 2

[0035] Eight 6-week-old SD strain rats were reared for 3 weeks on each feed shown in Table 3. The change in body weight during the experiment and the change in the degree of body fat on Day 21 are shown in Table 4. Further, triglyceride levels in serum after the animals were sacrificed are shown in Table 5.

[0036] As compared with other oil and/or fat feed groups, the oil and/or fat 1 feed group is significantly low in triglyceride levels in serum. It is understood that because the degree of body fat also indicates a similar result, accumulation of the fat in internal organs and adipose tissues is low given the feed using oil and/or fat 1.

Example 3

[0037] Eight 6-week-old SD strain rats were reared for 3 weeks on each feed shown in Table 6. The rats were allowed to run on a tread mill for 30 minutes once a day from the 14th day of rearing.

[0038] The change in body weight during the experiment, the change in the degree of body fat on Day 21, and the duration of running motility on the tread mill are shown in Table 7. Further, triglyceride levels in serum after the animals

were sacrificed are shown in Table 8.

Table 1

	Oil and/or fat preparation A	Oil and/or fat preparation B	Oil and/or fat preparation C	Oil and/or fat preparation D	Oil and/or fat preparation E	Rapeseed oil	MCT
Triglyceride	15%	14%	19.1%	19.2%	14.3%	98%	100%
Diglyceride	85%	86%	80.8%	80.8%	84.6%	2%	tr
Monoglyceride	tr	Tr	tr	tr	1.1%	tr	tr
Caproic acid	tr	Tr	tr	tr	37	tr	1
Caprylic acid	tr	Tr	tr	tr	tr	tr	1
Capric acid	100	Tr	27	29	63	tr	99
Palmitic acid	tr	4	tr	71	tr	4	tr
Stearic acid	tr	2	tr	tr	tr	2	tr
Oleic acid	tr	59	73	tr	tr	59	tr
Linoleic acid	tr	20	tr	tr	tr	21	tr
Linolenic acid	tr	10	tr	tr	tr	9	tr

Table 2

	Oil and/or fat preparation A	Oil and/or fat preparation B	Oil and/or fat preparation C	Oil and/or fat preparation D
CDM	60 hr or more	5.2hr	8.1hr	60 hr or more
Dissolution in mouth	excellent	excellent (liquid)	excellent	poor

Table 3

Oil and/or fat (one of the 4 kinds described below)	10 %
Casein	20 %
Mineral	3.5 %
Cellulose	4 %
Vitamin	1 %
Potato starch	61.5 %
Total	100.0 %

Oil and/or fat 1: Oil and/or fat preparation A

Oil and/or fat 2: Oil and/or fat preparation B

Oil and/or fat 3: Oil and/or fat preparation C

Oil and/or fat 4: Rapeseed oil

Table 4

Rearing period	Oil and/or fat	Oil and/or fat 2	Oil and/or fat 3	Oil and/or fat 4
Day 0 (body weight: g)	213.4±3.5	213.0±6.7	214.2±2.9	211.8±3.9
Day 14 (body weight: g)	311.8±6.1	355.2±9.0	341.5±3.6	371.0±5.5
Day 21 (body weight: g)	337.2±6.2	385.4±8.5	363.0±7.1	400.1±9.2
Degree of body fat (%: Day 21)	9.3±1.1	15.5±2.1	12.5±1.1	18.4±1.7

Table 5

Rearing period of 21 days	Oil and/or fat 1	Oil and/or fat 2	Oil and/or fat 3	Oil and/or fat 4
Serum triglyceride mg/ml serum	0.72±0.25	1.40±0.21	1.21±0.19	2.21±0.14

Table 6

Oil and/or fat (one of the 4 kinds described below)	10 %
---	------

Table 6 (continued)

Casein	20 %
Mineral	3.5 %
Cellulose	4 %
Vitamin	1 %
Potato starch	61.5 %
Total	100.0 %

Oil and/or fat 5: Oil and/or fat composition consisting of 59 % oil and/or fat preparation A and 41 % rapeseed oil (50 % capric acid diglyceride).

Oil and/or fat 6: Oil and/or fat composition consisting of 58 % oil and/or fat preparation B and 42 % rapeseed oil (50 % diglyceride derived from rapeseed aliphatic acid).

Oil and/or fat 7: Oil and/or fat preparation C (50 % capric acid-oleic acid diglyceride).

Oil and/or fat 8: Oil and/or fat composition consisting of 50 % MCT and 50 % rapeseed oil (50 % capric acid triglyceride).

Table 7

Rearing period	Oil and/or fat 5	Oil and/or fat 6	Oil and/or fat 7	Oil and/or fat 8
Day 0 (body weight: g)	210.0±3.9	211.5±4.2	211.8±3.0	212.5±6.8
Day 14 (body weight: g)	323.0±6.5	351.2±5.3	339.1±4.3	345.0±8.9
Day 21 (body weight: g)	348.0±6.1	375.3±9.6	359.0±6.7	368.5±11.5
Degree of body fat (%: Day 21)	12.8±1.1	18.9±1.2	14.0±0.6	15.3±0.4
Duration of motility (min)	41.0±2.9	32.3±3.2	36.5±1.5	28.5±2.1

Table 8

Rearing period of 21 days	Oil and/or fat 5	Oil and/or fat 6	Oil and/or fat 7	Oil and/or fat 8
Serum triglyceride mg/ml serum	0.76±0.09	1.43±0.20	0.96±0.09	1.26±0.10

[0039] From the results described above, the duration of motility in the oil and/or fat 5 feed group was improved as compared with the other oil and/or fat feed groups. Further, the oil and/or fat 5 feed group indicated lower degrees of body fat than in the other oil and/or fat feed groups. From the foregoing, it was found that the feed using the oil and/or fat of the present invention is converted very easily into energy and hardly accumulated as body fat.

Example 4

[0040] Now, an example of application to drink is illustrated. Oil and/or fat preparation A with the formulation shown in Table 9 was emulsified to prepare a drink.

[0041] As a result, it was found that its emulsification stability is high and the drink can be used without any particular problem with the flavor.

Example 5

[0042] Now, an example of application to a wheat flour product is illustrated. Oil and/or fat preparation A with the formulation shown in Table 10 was baked in a usual manner to produce cookies.

[0043] As a result, it was found that the cookies are excellent in flavor and also in releasability after baking, so the oil

and/or fat can be used as an equivalent to conventional oils and/or fats.

Example 6

5 [0044] Now, an example of application to whip cream is shown. An oily phase (oily ingredient liquid) and an aqueous phase (aqueous ingredient liquid) were prepared respectively according to the formulation shown in Table 11, mixed and preliminarily emulsified. Then, the resulting preliminarily mixture was homogenized at 60°C in a homogenizer. Then, the resulting emulsion was sterilized in an UHT sterilizer (145 °C, 3 seconds, Iwai Kikai Kogyo Co., Ltd.) and aseptically homogenized again at 70 °C in the homogenizer. The resulting emulsion was cooled to 15 °C, followed by
10 filling and subsequent aging for 1 day. The resulting emulsion was further whipped in a vertical whip machine (Kanto Kongoki Kogyo Co., Ltd.), to prepare a whip cream.

[0045] As a result, it was found that the whip cream is excellent in shape retention, flavor and eating-feeling in the mouth and the oil and/or fat can be used as an equivalent to conventional oil and/or fats.

Table 9

Ingredients	Ratio in food composition (%)
Oil and/or fat preparation A	20.0
Powdered skim milk	3.5
Protein (casein)	3.5
Yolk lecithin	0.7
Fructose	10.0
L-valine	0.5
L-leucine	0.5
L-isoleucine	0.5
L-arginine	0.5
Citric acid	0.1
Ascorbic acid	0.1
Perfume	0.1
Water	60.0
Total	100.0

Table 10

Ingredient	Weight in food composition (g)
Oil and/or fat preparation A	15.0
Corn starch	20.0
Wheat flour	50.0
Butter	5.0
Fructose	15.0
Common salt	0.5
L-valine	5.0
L-leucine	5.0

Table 10 (continued)

Ingredient	Weight in food composition (g)
L-isoleucine	5.0
L-arginine	5.0
Sodium bicarbonate	0.5
Water	10.0

Example 7

[0046] Now, an example of application to a cocoa drink is illustrated. The composition shown in Table 12 was mixed and homogenized at 70 °C in a homogenizer to prepare a cocoa drink.

[0047] As a result, it was found that the cocoa drink tastes suitably bitter, and there is no particular problem with its emulsified state, and the oil and/or fat can be used as an equivalent to conventional oils and/or fats.

Example 8

[0048] A commercial gelatin capsule No. 2 (weight: 53.3 mg) was charged with 150 mg of oil and/or fat composition A or E, to prepare hard capsules in a usual manner.

[0049] Hard capsule shells were charged in a usual manner with 500 mg of oil and/or fat composition A or E, to prepare hard capsules.

Table 11

	Ingredients	Weight ratio in food composition
Oily phase	Oil and/or fat preparation A	4.8
	Palm kernel hydrogenated oil (IV=2)	17.2
	Stearic acid monoglyceride (T-95, Kao Corporation)	0.1
	Oleic acid monoglyceride (O-95R, Kao Corporation)	0.1
Aqueous phase	Water	15
	Hydrous crystalline glucose (Showa Sangyo Co., Ltd.)	10
	Sunmalto S (Hayashibara Co., Ltd.)	20
	Oligotose (liquid) (Sanwa Denpun Kogyo Co., Ltd.)	28
	Powdered skim milk (3 % water content, 50 % lactose)	4.5
	Sucrose fatty acid ester (DK ester F-110, Daiichi Kogyo Seiyaku)	0.2
	Sodium hexametaphosphate	0.1

Table 12

Ingredients	Ratio in food composition (%)
Oil and/or fat preparation E	4.8
Powdered skim milk	3.5
Cocoa powder	1.1

Table 12 (continued)

Ingredients	Ratio in food composition (%)
Sugar	12.1
Corn syrup	1.5
Caramel	0.6
Chocolate essence	0.4
Gelatin	0.2
Water	75.8

Table 13

Hard Capsules	
Gelatin	70.0 %
Glycerol	22.9 %
Methyl paraoxybenzoate	0.15 %
Propyl paraoxybenzoate	0.15 %
Water	suitable amount
Total	100 %

Claims

1. A food comprising at least 0.5 % by weight of a partial glyceride derived from a C₂₋₁₀ aliphatic acid.
2. The food as claimed in Claim 1 wherein the number of carbon atoms of the aliphatic acid is 8 to 10.
3. The food as claimed in Claim 1 wherein the partial glyceride is a diglyceride.
4. The food as claimed in Claim 1 wherein the partial glyceride is a diglyceride, and at least one of the two aliphatic acids constituting the diglyceride is a C₈₋₁₀ aliphatic acid.
5. The food as claimed in Claim 1 wherein the partial glyceride is a diglyceride, and at least one of the two aliphatic acids constituting the diglyceride is a C₁₀ aliphatic acid.
6. The food as claimed in Claim 1 wherein the partial glyceride is a diglyceride, and both of the two aliphatic acids constituting the diglyceride are C₈₋₁₀ aliphatic acids.
7. The food as claimed in Claim 1 wherein the partial glyceride is capric acid diglyceride.
8. The food as claimed in Claim 1 comprising 0.5 to 85% by weight of the partial glyceride.
9. The food as claimed in Claim 1 wherein the partial glyceride is a diglyceride, at least one of the two aliphatic acids constituting the diglyceride is a C₁₀ aliphatic acid, and at least 4.0 % by weight of this partial glyceride is contained.
10. The food as claimed in Claim 1 or 9 wherein the food is a drink or an oral administration agent.
11. A method of storing glycogen in the body of a person who has ingested the foods of Claim 1.

Shaw + Co. 11/6/07
Sehach
Not a method
propyl paraoxybenzoate
clinical trial
action must
use is
similar
RT

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/00926

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.⁴ A23D7/00, 9/00, A23G3/00, A23C13/14, A23L2/38, A61K31/23

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.⁴ A23D7/00, 9/00, A23G3/00, A23C13/14, A23L2/38, A61K31/23

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP, 50-18001, B1 (Unilever N.V.), June 25, 1975 (25. 06. 75) & US, 3658555, A & DE, 1692541, A	1-9, 11
X	JP, 8-269478, A (Kao Corp.), October 15, 1996 (15. 10. 96) (Family: none)	1-11
A	JP, 8-311483, A (Loders Croklaan B.V.), November 26, 1996 (26. 11. 96) & EP, 739592, A	1-11
A	JP, 8-60180, A (Kao Corp.), March 5, 1996 (05. 03. 96) (Family: none)	1-11
A	JP, 4-300826, A (Kao Corp.), October 23, 1992 (23. 10. 92) (Family: none)	1-11

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
---	--

Date of the actual completion of the international search
March 30, 1998 (30. 03. 98)Date of mailing of the international search report
April 7, 1998 (07. 04. 98)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)